

When you have commenced treatment with medication and your diarrhoea has stopped for at least 48 hours then isolation precautions in the hospital may be ceased for you.

Your room or bed space should be cleaned each day. If this doesn't occur please inform the nurse caring for you.

### What happens when I go home?

There is no need to do anything extra at home. Good general personal and household hygiene including hand washing is important to prevent the spread of CDAD to others.

Eating utensils and dishes can be washed as normal.

You can return to work when you are well and your doctor says it is OK.

If you are a healthcare worker, check with Infection Prevention and Control before returning to work.

If you go to your GP or another healthcare facility please tell them you have had CDAD.

**If you have any questions about CDAD that have not been answered here, please ask your doctor or call Infection Prevention and Control.**

## Contact

### Infection Prevention and Control Health Directorate

Canberra Hospital And Health Services  
Po Box 11,  
Woden 2606 ACT  
Phone: 02 6244 3695  
Fax: 02 6244 4646

#### Accessibility

The ACT Government is committed to making its information, services, events and venues, accessible to as many people as possible.

- If you have difficulty reading a standard printed document and would like to receive this publication in an alternative format—such as large print or audio—please telephone 13 2281 or email HealthACT@act.gov.au.
- If English is not your first language and you require the translating and interpreting service—please telephone 131 450.
- If you are deaf or hearing impaired and require the TTY typewriter service—please telephone 13 3677, then ask for 13 2281.
- Speak and listen users—phone 1300 555 727 then ask for 13 2281.
- Internet Relay Users—connect to the NRS, then ask for 13 2281.

© Australian Capital Territory, Canberra, August 2013

This work is copyright. Apart from any use as permitted under the *Copyright Act 1968*, no part may be reproduced by any process without written permission from the Territory Records Office, Community and Infrastructure Services, Territory and Municipal Services, ACT Government, GPO Box 158, Canberra City ACT 2601.

Enquiries about this publication should be directed to ACT Government Health Directorate, Communications and Marketing Unit, GPO Box 825 Canberra City ACT 2601 or email: HealthACT@act.gov.au

www.health.act.gov.au | www.act.gov.au

Enquiries: Canberra 13ACT1 or 132281 | Publication No XX

# Clostridium difficile Associated Diarrhoea (CDAD)

## A Guide for Patients



## What is Clostridium difficile (C. difficile)?

Clostridium difficile (C. difficile) is a bacterium that can cause mild to very severe diarrhoea.

C. difficile is the most frequent cause of infectious diarrhoea in Australian hospitals and long-term care facilities. (Nursing homes etc)

C. difficile can cause complications such as pseudomembranous colitis (or life-threatening inflammation of the colon).

## How do people get Clostridium difficile Associated Diarrhoea (CDAD)?

Antibiotics alter the normal levels of good bacteria found in the bowels. When antibiotics destroy a person's good bowel bacteria, C. difficile bacteria can grow and produce toxins, which can damage the bowel and cause diarrhoea. This is known as C. difficile associated diarrhoea (CDAD). However, some people can have C. difficile bacteria present in their bowels and not show symptoms.

C. difficile bacteria and their spores are found in faeces. For healthy people, C. difficile does not pose a health risk. The elderly, those with other illnesses or who are taking antibiotics, are at a greater risk of infection.

People can get infected if they touch surfaces contaminated with C. difficile then touch their mouths. It is possible for healthcare workers to spread the bacteria to their patients if healthcare workers' hands are contaminated.

## How is CDAD diagnosed?

A sample of faeces from someone with diarrhoea is sent to the laboratory for testing. The laboratory checks for toxin-producing C. difficile bacteria. When C. difficile toxins are detected the patient is diagnosed as having CDAD.

## What are the symptoms of CDAD?

Symptoms include watery diarrhoea, fever, loss of appetite, nausea, and abdominal pain and tenderness.

## What can be done to prevent the spread of C. difficile?

In healthcare settings C. difficile infections can be limited through careful use of antibiotics and strict adherence to infection prevention and control measures.

Frequent hand hygiene is an effective way of preventing the transmission of healthcare associated infections. Hand washing with antimicrobial handwash is important during C. difficile outbreaks and is one of the best defences against further spread of the bacteria.

If you do not have access to hand washing facilities, frequent use of alcohol-based hand rub is encouraged. Most healthcare facilities provide alcohol-based hand rub at entrances to their buildings. Be sure to use them, but be aware that in CDAD they are less effective than hand washing as they do not destroy C. difficile spores.

If you work in or visit a hospital or long-term healthcare facility, wash your hands especially after using the toilet. Gloves should be worn by staff when caring for a patient with C. difficile infection or if in contact with his/her environment. Gloves should be changed between patients and hands should be washed after removing gloves.

When any antibiotic is prescribed for CDAD, follow your doctor, pharmacist, or healthcare provider's instructions and the directions on the label. Keep taking the antibiotic as prescribed in order to kill all of the C. difficile bacteria. If you have concerns about C. difficile and the medication you are currently using, talk to your doctor, pharmacist, or healthcare provider.

## What will happen while you are in hospital?

If you have CDAD you will be placed in a single room or in a shared room with other CDAD patients. All staff need to wear a yellow gown and gloves when entering your room or bed space. This is to prevent the spread of CDAD to other patients.

If you have CDAD the doctor will treat you with antibiotics that can kill the C. difficile bacteria and treat CDAD. It is important that you remain in your room to reduce the risk of spreading the C. difficile bacteria.

If you need to leave your room, your hands should be washed prior to leaving. It is important that you do not visit patients in the ward or in other wards in the hospital. (Parents of infants in the hospital nursery may visit their children. Please inform your nurses, wash your hands and wear a yellow gown when in the nursery.)

If you are required to visit other departments for specific tests e.g. X-ray, the staff caring for you will inform the department before you are transferred there. Staff such as wardsmen and the staff performing a procedure on you will wear a gown and gloves. If they do not, please inform them that you have C. difficile infection and they should be taking precautions.