

# CRYPTOSPORIDIOSIS

## What is cryptosporidiosis?

Cryptosporidiosis is an infection caused by a parasite called *Cryptosporidium*.

## What are the symptoms?

The symptoms usually include watery diarrhoea and abdominal pain. Fever, nausea, vomiting, and loss of appetite may also occur

Symptoms usually start from 1 to 12 days (average 7 days) after a person becomes infected. Symptoms may come and go and can last from days to weeks. Symptoms may be severe and longer lasting in people with weakened immune systems.

## How is cryptosporidiosis spread?

The *Cryptosporidium* parasite is found in the faeces of infected humans or animals.

People become unwell after swallowing the parasite. Infection can occur through

- person-to-person contact, particularly from young children to other children or their carers
- swallowing contaminated water from rivers, streams, springs, ponds, lakes, swimming pools, or the sea
- handling infected animals or their manure
- eating undercooked food, fruit or vegetables that have been contaminated or washed with contaminated water.

While people are most infectious when they have diarrhoea, they can still pass the infection on to others for several weeks after their symptoms have resolved.

## What treatment is available?

Most people usually recover with rest and extra fluids. Medication is generally not recommended, but may be prescribed by doctors in complicated cases.

## Who is at risk?

*Cryptosporidium* can infect anyone. However those who are most likely to become infected

with *Cryptosporidium* include:

- people in close contact with others who have cryptosporidiosis
- children who attend child care
- health care and child care workers
- swimmers who swallow even small amounts of swimming pool water
- people who drink untreated water
- international travellers, particularly to or from developing areas
- people who work with animals
- men who have sex with men

## How is it prevented?

### **Hand washing**

Hand washing is a very effective way of preventing person to person spread of *Cryptosporidium*. Hands should be washed with soap and running water for between 10 to 15 seconds:

- after going to the toilet or changing nappies
- after handling animals or their manure
- after working in the garden
- before preparing food or drinks
- before eating.

### **Nappy changing**

Nappies should not be changed on tables or counters where food is prepared or eaten. Change areas should be cleaned with warm soapy water and disinfectant after every nappy change.

### **Do not drink unpasteurised milk or untreated water**

People should avoid drinking unpasteurised milk or untreated water (for example from rivers, creeks, lakes, or dams). If untreated water is the only available source, boiling water will kill *Cryptosporidium*.

## Safe food handling

Make sure all foods are cooked thoroughly and wash or peel all raw vegetables and fruits before eating.

## How is it controlled?

Children who have diarrhoea should be kept home from preschool, childcare or play group until 24 hours after the diarrhoea has completely stopped.

People who work in food preparation, child care centres, aged care facilities, or health care facilities should not return to work until 48 hours after their diarrhoea has stopped.

People with cryptosporidiosis should not swim, wade, or paddle in swimming pools or recreational water facilities for at least two weeks after diarrhoea has stopped. Also, towels and linen should not be shared for at least two weeks after diarrhoea has stopped.

## Need more information?

For more information about salmonellosis, contact your doctor or call the Health Protection Service, Communicable Disease Control Information Line during business hours on **(02) 6205 2155**.

## Communicable Disease Control Section at Health Protection Service is

*responsible for the investigation and surveillance of notifiable or infectious conditions in the ACT in order to control or prevent their spread in the community. This includes the promotion of immunisation, education and other strategies that help to limit the spread of diseases.*

*Salmonellosis is a notifiable disease. Cases notified to ACT Health are investigated by Public Health Officers.*

## Acknowledgement

1. Heymann DL, 2015, *Control of Communicable Diseases Manual*, 20th edition.

## Accessibility

The ACT Government is committed to making its information, services, events and venues, accessible to as many people as possible.

- If you have difficulty reading a standard printed document and would like to receive this publication in an alternative format—such as large print or audio—please telephone 13 2281 or email HealthACT@act.gov.au.

- If English is not your first language and you require the translating and interpreting service—please telephone 131 450.
- If you are deaf or hearing impaired and require the TTY typewriter service—please telephone (02) 13 3677, then ask for 13 2281.
- Speak and listen users—phone 1300 555 727 then ask for 13 2281.
- Internet Relay Users—connect to the NRS, then ask for 13 2281.

© Australian Capital Territory, Canberra.

## Updated May 2017

This work is copyright. Apart from any use as permitted under the *Copyright Act 1968*, no part may be reproduced by any process without written permission from the Territory Records Office, Community and Infrastructure Services, Territory and Municipal Services, ACT Government, GPO Box 158, Canberra City ACT 2601.

Enquiries about this publication should be directed to ACT Government Health Directorate, Communications and Marketing Unit, GPO Box 825 Canberra City ACT 2601 or email: HealthACT@act.gov.au

[www.health.act.gov.au](http://www.health.act.gov.au) | [www.act.gov.au](http://www.act.gov.au)

Enquiries: Canberra 13ACT1 or 132281