

Want more information?

Contact the Health Protection Service,
Immunisation Unit

(02) 5124 9800 or

Winnunga Nimmityjah Aboriginal
Health Service

(02) 6284 6222

Visit our website at

<http://www.health.act.gov.au/our-services/immunisation>



Accessibility

If you have difficulty reading a standard printed document and would like an alternative format, please phone 13 22 81.



If English is not your first language and you need the Translating and Interpreting Service (TIS), please call 13 14 50.

For further accessibility information, visit:

www.health.act.gov.au/accessibility

www.health.act.gov.au | Phone: 132281

Publication No XXXXX

© Australian Capital Territory, Canberra December 2018

Immunise our children to stay healthy

Health Protection Service



ACT
Government

ACT Health

health.act.gov.au

Where can we go to get our child immunised?

Your doctor, Winnunga Nimmityjah or ACT Health Early Childhood Immunisation Clinic (02) 5124 9977 can immunise your child. Be sure to tell the doctor or nurse if your child is Aboriginal and/or Torres Strait Islander as they may need extra immunisations.

Why do babies need to be immunised at a young age?

Babies need help to fight disease. Immunisation protects against many diseases. You can help your child stay healthy by giving all immunisations on time.

How many immunisations will our child need?

At the following ages your child will need:

- 2 months – two injections and a liquid to swallow
- 4 months – two injections and a liquid to swallow
- 6 months – one injection
- 12 months – three injections
- 18 months – three injections
- 4 years – one injection

These immunisations will protect your child against 13 different diseases that can cause some types of pneumonia, diarrhoea, ear infections, whooping cough, measles and liver infections.

Your child might need extra immunisations if they have a chronic disease or if they were born early.

Annual influenza vaccination is also recommended for all Aboriginal and/or Torres Strait Islander children from 6 months of age.

Your child will be offered more immunisations when they get to high school.

When should our child start having their immunisation?

Your child can have their first immunisation at 6 weeks. The next immunisations should be given on time at 4, 6, 12 and 18 months of age and at 4 years.

What happens if we are late in taking our child for their immunisations?

To give your child the best protection, all immunisations should be given on time.

If you are late, your child can still have most of their immunisations but may miss out on receiving some that can only be given under a certain age.

Your doctor or nurse can help you catch your child up on any missed immunisations.

What can we expect after our child has been immunised?

Immunisation is safer than having the disease. Sometimes immunisation can cause minor and short-lasting reactions. These reactions could be mild fever (less than 38.5 degrees) or soreness, redness, itching and swelling around the injection site. Your child could also be grizzly and unsettled. These reactions usually last for 1–2 days.

If you are worried about your child after they have had their immunisations you should seek medical advice from your doctor, hospital, Maternal and Child Health Nurse or Winnunga Nimmityjah Aboriginal Health Service. You can also contact the Health Protection Service, Immunisation Unit on (02) 5124 9800 from Monday to Friday between 8.30 am and 4.30 pm.

